

University of Groningen

Assessing the performance of forthcoming Infrared telescopes

Bisigello, Laura

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Bisigello, L. (2017). *Assessing the performance of forthcoming Infrared telescopes*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

PROPOSITIONS

accompanying the dissertation

ASSESSING THE PERFORMANCE OF FORTHCOMING INFRARED TELESCOPES

1. In a microwave kinetic inductance detector array, the responsivity based on the readout frequency response is inversely proportional to the responsivity based on a calibrated optical source, for a small difference in optical power (Chapter 2).
2. By using a two-dimensional Gaussian function, it is possible to characterise the level of crosstalk on microwave kinetic inductance detector arrays and correct images for this effect a posteriori (Chapter 3).
3. At $z < 7$, having ancillary *Hubble Space Telescope* data to complement the NIRCam broad-band observations obtained with the *James Webb Space Telescope* is fundamental to limit the fraction of outliers in redshift (Chapter 4).
4. When observing galaxies with the two imaging cameras on the *James Webb Space Telescope*, observations with the MIRI/F560W and/or F770W broad-band filters are essential to have a good estimation of stellar masses at $z = 10$ (Chapter 5).
5. The presence of numerous nebular emission lines makes the photometric redshift and stellar mass estimation challenging (Chapter 4, 5).
6. The multi-Gaussian decomposition of the specific star-formation-rate distribution allows for the derivation of the main-sequence of star-forming galaxies without selection effects (Chapter 6).
7. The fraction of starbursts increases with redshift and decreases with stellar mass (Chapter 6).
8. A high percentage of PhD students and post-docs leave astronomy because of the nomadic lifestyle associated with the astronomy career.
9. North Italy is not Milan.
10. It is possible to obtain a PhD degree in astrophysics without working during weekends.
11. Astronomical observations are based on the principle of the “maximum results with the minimum effort”.

Laura Bisigello